Going rogue: In search of passionate and unencumbered talk about energy in the blogosphere.

Lane H. Seeley, Eleanor W. Close, Lezlie S. DeWater,

Rachel E. Scherr

AAPT Summer Meeting, Portland, OR, July 21, 2010

ENERGYPROJECT







The SPU Energy Project




Dedicated to the exploration of ideas about energy

Energy on the Web?






























Energy we learn about

Energy we care about





















Experimental Approach 'Study people in their natural habitat'







'Gaussian Gun Energy Test'

futuristic
@www.alter.si








'Gaussian Gun Energy Test'

futuristic @www.alter.si



















Iratcliff - 'Can you make a channel so the balls fall back down and feeds back in so that it can repeat continuously? And, if so, what is keeping you from putting a little windmill contraption as the ball shoots out to convert a little bit of the linear energy into rotational energy? If you could do this, wouldn't that qualify as a 'free-energy' device, as you would harvest a bit of the energy gain?'



Frenkystromar — 'I could bring back ball with no problem. But there is already one ball attracted to the magnet. So first this ball should be removed and then the effect could be repeated. But it takes a lot of energy to remove the ball that was attracted in the first round. The big question is: Is the kinetic energy of the ball that rolls away bigger than the energy needed to remove the ball that was attracted to the magnet?'

Jratcliff - 'Can you make a channel so the balls fall back down and feeds back in so that it can repeat continously? And, if so, what is keeping you from putting a little windmill contraption as the ball shoots out to convert a little bit of the linear energy into rotational energy? If you could do this, wouldn't that qualify as a 'free-energy' device, as you would harvest a bit of the energy gain?'

Creative Respect Respect for Refined

Insight Evidence Inquiry

Frenkystromar — 'I could bring back ball with no problem. But there is already one ball attracted to the magnet. So first this ball should be removed and then the effect could be repeated. But it takes a lot of energy to remove the ball that was attracted in the first round. The big question is: Is the kinetic energy of the ball that rolls away bigger than the energy needed to remove the ball that was attracted to the magnet?'

A Rich Resource

The blogosphere provides a rich, searchable and relatively untapped database of thinking about energy in which 'students':

- •Negotiate their own language for discourse (free-energy, overunity, HHO, etc...)
- Talk about what they care about
- •Demonstrate a capacity for autonomous productive disciplinary engagement



A Useful Resource

We can use this resource to:

- •Identify questions which will engage student curiosity and prior knowledge
- Anticipate the landscape of student thinking about these questions
- •Help students construct a bridge between the energy they learn about and the energy they care about





Seattle Pacific University

Physics Education Research Group



Faculty Members:

Hunter Close

Eleanor Close

Lane Seeley

Stamatis Vokos

Master Teacher:

Lezlie DeWater

Researchers:

Rachel Scherr

Sam McKagan

Project Coordinator:

Julie Glavic

High School Intern:

Emma Kahle





Personal Engagement on the Web







Personal Engagement on the Web

Gravity Powered Plane




You can never have greater than 100% efficiency


Its called over unity and YES it is absolutely possible and has been demonstrated many times.

Heat Pumps have a CoP 3.0 (coefficient of performance).
One times electric power in, three times the heat out. (PulseFuelNerd)















Theoretical Approach 'Real World Problems'

